



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Biopesticides and Pollution Prevention Division (7511P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

71038-6

Date of Issuance:

7/22/2016

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Fortune Aza 6% WP

Name and Address of Registrant (include ZIP Code):

Fortune Biotech, Ltd.  
6-6-125 ANNAM GARDENS, KAVADIGUDA  
SECUNDERABAD, 500 380  
INDIA

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Make the following labeling change before you release this product for shipment:  
Revise the EPA Registration Number to read, "EPA Reg. No. 71038-6".
3. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Signature of Approving Official:

Andrew Bryceland, Team Leader  
Biochemical Pesticides Branch  
Biopesticides and Pollution Prevention Division (7511P)  
Office of Pesticide Programs

Date:

7/22/2016

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 4/5/16.

If you have any questions, please contact Chris Pfeifer of my team by phone at 703-308-0031 or via email at [pfeifer.chris@epa.gov](mailto:pfeifer.chris@epa.gov).

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive, with the first name being the most prominent.

Andrew Bryceland, Team Leader  
Biochemical Pesticides Branch  
Biopesticides and Pollution  
Prevention Division (7511P)  
Office of Pesticide Programs


Enclosure

# Fortune Aza 6% WP

Master Label

This product is botanically derived from the Neem plant "*Azadirachta Indica*" Azadirachtin, a botanical insecticide.

**Sublabel A: For Agricultural and Commercial Use**  
**Sublabel B: For Residential Use**



**ACTIVE INGREDIENT:** Azadirachtin ..... 6.0%  
**OTHER INGREDIENTS:** ..... 94.0%  
**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

EPA Registration Number: 71038-A  
EPA Establishment No. : 71038-IND-1

Manufactured By:  
**FORTUNE BIOTECH LIMITED**  
6-6-125, Annam Gardens, Kavadiguda  
Secunderabad, 500380 India

US Contact Address:  
6860 N. Dallas Parkway, Suite 200  
Plano, TX 75024

Net Contents: \_\_\_\_\_

## Sub Label A

This sub-label bears direction agricultural/commercial use with specific application rates associated with the package size and area to be treated.

# Fortune Aza 6% WP

This product is botanically derived from the Neem plant "Azadirachta Indica" Azadiractin, a botanical insecticide.

### [Product Features]

- **Non-oil based, water soluble powder**
- **Effective on a very wide spectrum of insects**
- **Use as a spray, drench, tree injection or chemigation**
- **May be applied as directed to any food crop up to and including the day of harvest**

For use on turf grass, outdoor shrubs, trees and ornamentals

For ornamental greenhouse, shade house, interiors cape and nursery use

For mushroom house use

For use on outdoor food crops

For control, growth control, antifeedant and repelling Insects such as aphids, armyworms, beetles, budworms, cutworms, fungus gnats, leafhoppers, leafminers, leafrollers, lepidopterous larvae, loopers, mushroom flies, sawflies, thrips, webworms, whiteflies and other pests as listed; and plant parasitic nematodes such as dagger, golden, and root knot nematodes in vegetables, fruits, nuts, coconuts, agronomic crops and ornamental plants. For residential and commercial lawn, flowers and vegetable gardens, farms, forests, sod farms, nurseries, greenhouse-food and ornamental plants, mushrooms, nursery plants, interiorscapes, landscapes, turfgrasses and golf courses.

**ACTIVE INGREDIENT:** Azadirachtin ..... 6.0%

**OTHER INGREDIENTS:** ..... 94.0%

**TOTAL:** ..... 100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>Hotline Number</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies, phone 24 hours a day, National Pesticide Telecommunication Network at 1-800-858-7378.	

See Side/Back Panel for First Aid and Precautionary Statements

**EPA Registration Number: 71038-A**

**EPA Establishment. No.: 71038-IND-1**

**Net Contents:** \_\_\_\_\_

Manufactured By:  
FORTUNE BIOTECH LIMITED  
6-6-125, Annam Gardens, Kavadiguda  
Secunderabad, 500380 India

US Contact Address:  
6860 N. Dallas Parkway, Suite 200  
Plano, TX 75024

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical resistant gloves

Follow manufacturer's instructions for cleaning /maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any irrigation system unless the chemigation instructions on this label are followed. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical resistant gloves

### NON-AGRICULTURAL USE REQUIREMENTS

These requirements apply to uses of this product that are NOT within the WPS for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses including golf courses and other non-agricultural uses, do not enter treated areas without protective clothing until sprays have dried.

## **PRODUCT DESCRIPTION**

This product is a pale yellow/white, amorphous powder containing 6% by weight azadirachtin. It will instantly dissolve in water to give a solution that is ready for spray applications for pest control. Non-Oil based and highly effective as a powder.

Use this product for pre-harvest treatment of fruits and vegetables in case of sudden pest infestations. This product is effective on a very wide spectrum of insects and pests as listed on this label.

Use this product on a wide variety of plants as listed indoors and outdoors. If plans are made to use this product on plants not listed on this label, it is recommended that a small area such as a leaf, stem, or branch be “test sprayed”, first, and checked several days later to make sure that leaf wilting or damage does not occur.

When used as directed, this product will destroy targeted insect larvae when they, (1), eat sprayed plants, or (2), come in contact with the spray. This product eliminates insects by stopping the insect’s growth, and is effective on all insects listed, insect larval stages and pupae.

## **MODE OF ACTION**

This product controls insects in the larval, pupal, and nymphal stages by interfering with the metabolism of ecdysone. Insects typically die between larval to larval, larval to pupal, nymph to nymph molts, or during adult eclosion.

## **COMPATIBILITY**

This product has been found to be compatible with the most commonly used non-alkaline insecticides, fungicides and water soluble fertilizers in the neutral pH range. Check compatibility by using the correct proportion of each of the products application rate in a quart or gallon container. Solubilize this product first in the mixture. Test the tank-mix combinations for possible adverse effects (such as settling out, flocculation, etc.) and for phytotoxic effects on a small sample of plants prior to use. As environmental conditions can alter the interactions between compounds, test compatibility for both new and previously used combinations. Avoid mixtures of several materials and very concentrated spray mixtures.

Do not use this product with Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials. Use mildly alkaline mixtures immediately after mixing to prevent loss of insecticidal activity.

When using this product in combination with other products, use this product at the rate, or half the rate, specified in the Use Rate table. Follow the directions for use, precautions and limitations for use on all of the product labels used in the combination. Some suggested tank mix combinations are as follows:

Fortune Aza 6% WP plus endosulfan\*

Fortune Aza 6% WP plus chlorpyrifos\*

Fortune Aza 6% WP plus acephate\*

Fortune Aza 6% WP plus *Bacillus thuringiensis*\* (BT)

Fortune Aza 6% WP plus bifenthrin\*

Fortune Aza 6% WP plus esfenvalerate\*

Fortune Aza 6% WP plus abamectin\*

Fortune Aza 6% WP plus diflubenzuron\*

Fortune Aza 6% WP plus pyrethrum + piperonyl butoxide (for fogging use)\*

- Always follow the manufacturer’s Directions for Use and Precautionary Statements.

Use Fortune Aza 6% WP on vegetables, coconut palms and other food crops with such chemicals as Endosulfan.

## **APPLICATION INSTRUCTIONS**

This product is exempt from tolerances and may be applied as directed to any food crop up to and including the day of harvest at a rate not exceeding 0.75 lb (20 grams active ingredient) per acre per application.

## **READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE**

To apply this product select a suitable power or pump pressure type sprayer or a hand held trigger type sprayer that will deliver a forceful, fine, leaf and fruit covering, wetting, spray mist. To get thorough spray coverage on waxy or pubescent plant surfaces the addition of small amount of a suitable sticker agent (such as NuFilm P) added to the spray mix, at the recommended rates may give better foliage, insect coverage and control.

**APPLICATION METHOD AND EQUIPMENT:** Apply this product as a foliar spray or a drench to soil or soil-less media (e.g., greenhouses and mushroom houses) to control insects and nematodes. When needed, drench soil to control soil-borne pests, including soil-borne larvae of foliar insect pests. When applying as a drench, avoid excessive leaching. Apply this product through sub-surface soil treatment equipment (e.g. turf grass). To repel adult flies, apply through fogging equipment. Always follow equipment manufacturers use directions.

Apply this product by using any powered or manual pesticide application equipment, which includes but is not restricted to: high-volume, low- volume, ultra-low volume, electrostatic, fogging, and chemigation. Follow the manufacturer's instructions when using these types of equipment.

**For optimum results, 2 to 3 applications made at 7 to 10 day intervals is recommended, unless otherwise specified.** Foliar applications should be made to both sides of leaves. In addition, a sticker agent used as per the manufacturer's instructions may improve product performance.

**APPLICATION METHOD AND EQUIPMENT:** Apply this product as a foliar spray to control insects and nematodes. Apply this product through sub-surface soil treatment equipment (e.g. lawn grass). To repel adult flies, apply through fogging equipment. Always follow equipment manufacturers use directions.

Apply this product by using any powered or manual pesticide application equipment. Follow the manufacturer's instructions when using these types of equipment.

**For optimum results, 2 to 3 applications made at 7 to 10 day intervals is recommended, unless otherwise specified.** Foliar applications should be made to both sides of leaves. In addition, a sticker agent used as per the manufacturer's instructions may improve product performance.

### **USE RATE DIRECTIONS** **FOR KEY PESTS BY USE SITE**

This product is intended for use on outdoor plants and food crops, mushroom houses, plants grown indoors or in greenhouses, shade cloth, interiors capes and nurseries. It can be used to control any of the following insects and nematodes.

This product label rates specify dry ounce (weight/acre (high rate) and tsp or tbspc/1000 sq. ft. (low rate). These label rates provide a high and low dose application of this product.

<b>Abbreviation &amp; Conversion Table</b>	
tsp	teaspoon
tbsp	tablespoon
A	acre
3 tsp = 1 tbsp	
1 A = 43,560 sq. ft.	

**High Rate**

6oz/ 50 gal water / Acre (6oz. is approximately 170 tsp / 50 gal water /acre) (approximately 4 tsp / 1000 sq ft.)

**Low Rate**

1 tsp/ 1 gal water / 1000 sq. ft. (1 tsp. is approximately 44 tsp / 44 gal water / acre)

**COMMON USE CONDITIONS:**

**Low Rate** (Commonly used for preventative treatments before signs of infestation.)

1 tsp/ 1000 sq. ft.

**Medium Rate** (Used for most treatments. For preventative to medium infestations when pests are present.) 2-3 tsp/ 1000 sq. ft.

**High Rate** (Used for difficult to manage pests or high infestations.)

4 tsp/ 1000 sq. ft.

Use the tables below to determine the appropriate use rate for your site / pest combination.


**PRODUCT USE RATE DIRECTIONS**  
**FOR KEY PESTS BY USE SITE**

This product is intended for use on outdoor plants and food crops, mushroom houses, plants grown indoors or in greenhouses, shade cloth, interiors capes and nurseries. It can be used to control any of the following insects and nematodes.

Use the tables below to determine the appropriate use rate for your site / pest combination.

**PRODUCT PEST CONTROL CHART: USE RATES for indoor and outdoor plants including, FOOD CROPS, TREES, TURFGRASS, NURSERY, GREENHOUSE, INTERIORSCAPE, & LANDSCAPE PLANTS.**

<b>PEST</b>	<b>RATES: Fortune Aza 6% WP oz's./Acre-tsp./1,000 sq ft.</b>	<b>COMMENTS For Spray, Drench or Chemigation.</b>
<b>WHITEFLIES</b> , such as: Green-house whiteflies, Silverleaf white flies, Woolly whiteflies.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Make sure that spray covers upper, lower and all surfaces of leaves fruit and twigs.
<b>LEAF MINERS</b> , such as Azalea leafminers, Birch leafminers, Citrus leafminers, Serpentine leafminers	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Apply to new growth in spring before new larvae enter plant foliage. Repeat application at 10 to 14 day intervals if new infestations are expected.
<b>SCALE</b> , Crawlers: such as Brown Soft Scale, California red scale, Coffee Scale, Olive Scale, San Jose Scale.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Make sure to thoroughly spray upper, lower and all surfaces of leaves and twigs.
<b>MEALY BUGS</b> Such as Citrus Mealybugs	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray to thoroughly cover twigs and leaves.
<b>THRIPS</b> , such as: Citrus thrips, Onion thrips, thrips palmi,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray in spring when young nymphs first appear on foliage.
<b>APHIDS</b> , such as: Cotton aphids, Green peach aphids, Pea aphids, Potato aphids	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray to wet lower side of leaves when "leaf curl" first appears.
<b>PSYLLIDS</b> , such as: Pear psylla	Same as above	Spray for new "instar" nymphs appearing on new discolored foliage.
<b>BUGS</b> , Nymphs of: such as Box-elder bugs, Chinch bugs, lygus bugs, spittle bugs, stink bugs,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray early when nymphs are young. This product will control " <u>instar</u> " growth until they die.
<b>FLIES</b> , Larvae of: such as Blueberry Maggot, Cherry Maggot, Crane Flies, Fruit flies, Midges, Onion Maggots, Tip worms, Walnut husk fly larvae.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	For Food and Non-food crops spray when larvae first appear.
<b>SAWFLIES</b> , Larvae of: such as: European Pine Sawflies, Yellow headed pine sawflies,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray when first larvae appear when plants start new growth.
<b>CATERPILLARS</b> , Such as: Armyworms, Artichoke plume moth, Bagworms, Bollworms,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray when first larvae worms appear. Repeat applications in 7 to 10 days. For continued pest control in the spring

Budworms, Cabbage butterflies, Cabbage loopers, Cankerworms, Caseworms, Corn Earworms, Cutworms, Diamond back moths, Fireworms, Fruitworms, Grapeleaf skeletonizer, Gypsy moths, Hornworms, Imported Cabbage worm, leaf perforators, Leafrollers, Melonworms, Navel orangeworms, Oblique banded Leafrollers, Omnivorous Leafrollers, oriental fruit moths, Pickleworms, Pine tip moths, Pinworms, Red banded leaf rollers, Sod webworms, Soybean loopers, Tent Caterpillars, Tobacco budworms, Tussocks moth larvae.		or fall when insect infestations are expected spray ornamentals and other plants at intervals of 2 to 3 weeks.
<b>BEETLES</b> , Larvae of: such as Bark beetles, Blueberry Flea beetles, Boll weevils, Colorado potato beetles, Flea beetles, Japanese beetles, Leaf beetles, Mexican, bean beetles, Phylloxera, Rose Chafers, Twig girdlers	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray when pests first appear. For Food Crops. Repeat application after 7 to 10 days. <u>Do not use with oil!</u> Make sure that all plant surfaces are thoroughly spray treated. Repeat in 5 to 7 days if required.
<b>WEEVILS</b> , Such as Black vine weevils, Pepper weevils, Strawberry vine weevils.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Foliar anti-feedant sprays will stop adult feeding. Make at least 3 to 4 applications 10 days apart.
<b>BORERS</b> , Larvae such as: Peach twig borer, Peach tree borers, Cranberry borers,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Thoroughly spray in spring after egg hatch to control young larvae.
Emerald Ash Borer.	 <p>The diagram is a decision tree for Emerald Ash Borer (EAB) treatment. It starts with two main categories of areas: 'Within a quarantined county, a state quarantine generally-infested area or less than 15 mi. from a known EAB infestation' and 'In a county where EAB has not been detected and greater than 15 mi. from a known EAB infestation'. For the first category, there are two paths: 'EAB infested' and 'Suspicious EAB symptoms or No EAB symptoms'. For 'EAB infested', if there is '≥ 30% canopy thinning and/or dieback', treatment is not recommended; if '&lt; 30% canopy thinning and/or dieback', the treatment is 12.5 ml/in. dbh. For 'Suspicious EAB symptoms or No EAB symptoms', if there is '≥ 30% canopy thinning and/or dieback', treatment is 12.5 ml/in. dbh; if '&lt; 30% canopy thinning and/or dieback', treatment is 12.5 ml/in. dbh. For the second category, there are two paths: 'Suspicious EAB symptoms' and 'No EAB symptoms'. For 'Suspicious EAB symptoms', if there is '≥ 30% canopy thinning and/or dieback', treatment is 12.5 ml/in. dbh; if '&lt; 30% canopy thinning and/or dieback', treatment is 12.5 ml/in. dbh. For 'No EAB symptoms', if the tree is '&gt; 16 in. dbh', treatment is 10 ml/in. dbh; if '8 - 16 in. dbh', treatment is 8 ml/in. dbh; if '&lt; 8 in. dbh', treatment is 6 ml/in. dbh. A note at the bottom states: '* Tree condition may compromise treatment effectiveness'.</p>	
<b>MOLE CRICKETS</b> , nymphs and young "in-stars". Turf <u>Treatment.</u>	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	For turfgrass, spray to drench turf for young cricket nymphs in spring. Stops young from growth to adults.
<b>MUSHROOM FLIES</b> , Nematodes and Phorid Flies	Mix ½ oz. in 1 to 2 gallon of water and mist over, (or drench) 1,000 sq ft.	See "For Mushrooms" Section on this label.

## USE SITES

This product can be used on Green-house: food crops, such as: Brassica (cole) crops, cucurbits, eggplants, herbs and spices, legumes, peppers, tomatoes.

**MUSHROOMS**, Varieties such as: Agaricus, enoki, maitake, oyster, shitake, and other specialty mushrooms

**FOOD CROPS**, including: .....

**Root, and tuber vegetables**, such as: Artichoke, beets carrots, ginger, horseradish, potatoes, radishes, rutabagas, sweet potatoes, turmeric, turnips, yams.

**Leafy vegetables** (including Brassica leafy vegetables), such as: Amaranth, broccoli, Brussels sprouts, cabbage, cauliflower, celery, chervil, Chinese cabbage, collards, cress, endives, fennel, kale, kohlrabi, lettuce, mizuna ('mustard greens, parsley, purslane, rape greens, rhubarb, spinach, Swiss chard.

**Legume vegetables**, such as: beans (field, kidney etc.), chick-peas, cowpeas, guar, jackbeans, lablab beans, lentils, peas, pigeon peas, soybeans, sword beans.

**Fruiting vegetables**, such as: Eggplants, ground-cherries, pepinos, peppers, pimentos, tomatillos, tomatoes.

**Cucurbit vegetables**, such as: bitter melons, Chayotes, Chinese wax gourds, citron melons, cucumbers, gherkins, gourds, **muskmelons** (such as cantaloupes, casabas cranshaw etc.) pumpkins, squash, watermelons.

**Citrus fruits**, such as: Calamondins, citrus citrons, citrus hybrids, Grapefruits, Kumquats, Lemons, Limes, Mandarins, Oranges, pumellos, satasuma mandarins.

**Pome fruits**, such as: Apples, crabapples, loquats, mayhaws, oriental pears, pears, quinces.

**Stone fruits**, such as: Apricots, cherries, nectarines, peaches, plums, prunes.

**Berries**, such as: Blackberries, caneberries, blueberries, currants, cranberries, elderberries, gooseberries, huckleberries, loganberries, raspberries, strawberries, youngberries.

**Cereal grains**, such as: Barley, buckwheat, corn, millet, oats, popcorn, rice, rye, sorghum, teosinites, triticale hybrids, wheat, wild rice.

**Herbs and spices**, including but not limited to: allspice, ahgelica, anise, annatto, balm, basil, black and white peppers, borage, burnet, camomile, caper buds, cardamom, caraway, cassia, catnip, celery seeds, chervil, chives, cinnamon, caraway, cloves, coriander (cilantro), costmary, cumin, curry leaf, dills, fennels, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, mace, marigolds, marjoram, mustard seeds, nasturtium, nutmeg, parsley, pennyroyal, pepper (black & white), poppy seeds, rosemary, rue, saffron, sage, savory, sweet bay (bay leaf), tansy, tarragon, thyme, vanilla, wintergreen, woodruff, wormwood.

**Bulb vegetables** such as Garlic, leek, onions, shallots.

**Nuts**, such as: Almonds, beechnuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, Coconuts, filberts, hickorynuts, Imacadamia, pecans, pistachios, walnuts.

**Oilseed crops** such as: Canola, castor, crambe, guar, jojoba, peanuts, rape, safflower, sesame, soybean, sunflower.

**Tropical fruits**: such as Atemoyas, bananas, breadfruits, cherimoyas, durians, guavas, malangas, mangos, papayas, passionfruits, starfruits.

**Other food & non-food crops**: such as Asparagus, avocados, birdseed, cacao, coffee, edible flowers, feijoa, figs, ginseng, grapes, guayule, hops, kiwis, okras, olives, palms, papayas, pawpaws, persimmons, pineapples, rambutans, sugarcane, tamarillos, tea, tobacco, water chestnuts, watercress.

**Ornamental Plants**: such as: African violets, ageratum, aster, aucuba, begonia, cacti, calendula, calla, carnation, ceanothus, chrysanthemum, cineraria, coleus, cyclamen, daffodil, dahlia, delphinium, ficus, foliage plants, fuschia, gardenia, geranium, gloxinia, hyacinth, hydrangea, iris, ivy, lily, maidenhair fern, marigold, narcissus, orchid, pansy, pelargonium, peony, phlox, pittisporum, poinsettia, pyracantha, rubber plant, snapdragon, stock, tulip, wandering jew, yew, yucca, zinnia.

**Ornamental Trees and Shrubs**: such as: Andromeda, Arbovitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, camellia, cedar, chamaecyprus, dogwood, douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honeylocust, horsechestnut, ilex, juniper, larch laurel, lilac, linden, London plane, magnolia, manvillia, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, pine, phinota, pines, plane tree, poplar, privet, quince, rhododendron, roses, spruce, sycamore, white cedar and white pine.

## FOR TURF GRASSES

Bent grass	Fescue
Bermuda grass	Ryegrass; annual
Bluegrass annual & perennial	Ryegrass; perennial
Buffalo grass	Wheat grass
Centipede grass	Zoysia grass

For control of Sod Webworms, Cutworms, Aphids, Leafhoppers, ants, and chiggers: use a suitable pressure sprayer and mix 1 tbsp. in 2 to 3 gal. of water and apply to 2,500 sq. ft. of turf. Apply when insect larvae first appear and if necessary repeat application in 10 to 14 days. The use of an approved "spreader sticker" may help the spray to penetrate turf down to the larvae/worm feeding area.

#### **For Mushrooms and the Mushroom House**

For Mushroom Flies, Nematodes and Phorid Flies use this product at the rate indicated on the PEST CONTROL CHART as a drench to the casing layer, media or compost. Make 4 to 5 applications 7 to 10 days apart.

To repel fly adults, apply with fogging equipment at the first sign of activity. Can be applied between breaks up to the final flush.

### **Application by Trunk Injection**

#### **Directions for Use**

Inject into the trunk flare or within 36" of the soil level. Place the injection sites in the first few sapwood elements (growth rings). Drill holes using a clean sharp drill bit (brad point drill bits are recommended). Drill through the bark and into the sapwood.

#### **Calculating Application Rate**

The dosage and number of application sites are based on tree diameter (dbh). To determine the number of application sites and dose rate per tree:

1. **Determine the Tree Diameter (DBH):**

Measure the tree diameter in inches (or centimeters) at chest height [54" (135 cm) from the ground] to find the diameter at breast height (dbh). If measuring tree circumference, divide the circumference by 3 to obtain the dbh.

2. **Calculate # of Injection Sites:**

By Micro-injection (QUIK-jet®, Air/Hydraulic): Calculate the number of injection sites by dividing the dbh in inches by 2 (or cm dbh by 5). This is equivalent to one drill hole for every 6" (15 cm) of tree circumference. By Micro-infusion® (Tree I.V.): Calculate the number of injection sites by dividing the dbh in inches by 3 (or cm dbh by 7.5). This is equivalent to one drill hole for every 8" (20 cm) of tree circumference.

3. **Determine the Dose:**

Measure the amount of this product needed following the table: *Use Rate Instructions for Tree Injection*

4. **Determine Dose per Injection Site:**

Divide the total dose by the number of injection sites to determine the dosage per injection site

In resinous conifers (such as pine and spruce) you may inject each site shortly after drilling to avoid slow uptake on account of resin flow. In palms, only one injection site is generally required.

1. Locate the application site 1-3' (30-90 cm) from the soil level
2. Drill depth is 1/3 the total diameter or 4" (10 cm) deep into the stem (whichever is less).
3. Refer to table: *Use Rate Instructions for Palm Injection* for dosages to apply.

#### **Application Equipment**

This product may be used with tree injection devices that meet the label requirements. For all injection systems, read carefully and follow the manufacturer's direction for use.

#### **Mixing Procedures for Tree Injections**

Prepare the injection solution by mixing 8 level teaspoons (8 grams) of this product in 3.38 fl. oz. (100 mls) of water. Only mix the amount you plan to use so for smaller amounts refer to the table: *Use Rate Instructions for Tree Injection*.

#### **Application in Trees**

1. Inject 4 mls of solution every 6" (15 cm) of stem circumference in trees <8" dbh (20cm).
2. Inject 6 mls of solution every 6" (15 cm) of stem circumference in trees 8-16" dbh (15 – 40cm).
3. Inject 8 mls of solution every 6" (15 cm) of stem circumference in trees >16" dbh (>40cm).

**Use Rate Instructions for Tree Injection**

DBH"	cm DBH	level tsp. Product	Milliliters of water	ml/2.5 cm (inch) DBH	Average Number of injects	ml/inject
5	12.5	2	12	2	3	4
6	15.0	2	12	2	3	4
7	17.5	2	16	2	4	4
8	20.0	4	24	3	4	6
9	22.5	4	30	3	5	6
10	25.0	4	30	3	5	6
11	27.5	6	36	3	6	6
12	30.0	6	36	3	6	6
13	32.5	6	42	3	7	6
14	35.0	6	42	3	7	6
15	37.5	8	48	3	8	6
16	40.0	8	48	3	8	6
17	42.5	10	72	4	9	8
18	45.0	12	72	4	9	8
19	47.5	12	80	4	10	8
20	50.0	12	80	4	10	8
21	52.5	14	88	4	11	8
22	55.0	14	88	4	11	8
23	57.5	14	96	4	12	8
24	60.0	16	96	4	12	8
25	62.5	16	104	4	13	8

For Trees larger than 25" (62.5 cm) dbh apply 4 ml/2.5 cm (inch) dbh.

**Use Rate Instructions for Palm Injection**

Canopy or Tree Size	Tsp Product	Milliliters of water	Minimum Number of Injection Points Needed
Small Canopy or Tree	2	10	1
Medium Canopy or Tree	4	20	1
Large Canopy or Tree	6	30	1

Packet Size	Mylar Packet	Water Soluble Packet
0.07 oz. packets (approx. 2tsp)	not available	yes
0.75 oz. packets (approx. 22tsp)	yes	yes
3.00 oz. packets (approx. 85tsp)	yes	yes

## Dose/Rate Spray Applications

This product is measured in dry ounces (weight) and approximate teaspoons for each packet size. Packets come in various sizes. Depending on the type of pest and timing of treatment you may use low, medium, or high rates of application.

**NOTE: Low Rates** are commonly used for preventative treatments before signs of insects. **Medium Rates** are used for most treatments for preventative to medium infestations when pests are present. **High Rates** are used for difficult to manage pests or for heavy infestations.

### *Product packet directions for mixing, and dosing in spray applications.*

*Packet sizes are determined by using the most economical size and no partial packets.*

Sq Ft.	Low Rate		Medium Rate		High Rate	
	Product	Water	Product	Water	Product	Water
<b>1,000</b>			(1) 2tsp packet	1-2 gal	(2) 2tsp packets	1-4 gal
<b>5,000</b>	(3) 2tsp packets	5-10 gal	(5) 2tsp packets	5-10 gal	(1) 0.75 oz packet	10-20 gal
<b>10,000</b>	(5) 2tsp packets	10-20 gal	(1) 0.75oz packet	10-20 gal	(2) 0.75 oz packets	20-40 gal
<b>20,000</b>	(1) 0.75oz packet	20-40 gal	(2) 0.75 oz packets	20-40 gal	(1) 3.00 oz. packet	50-100 gal
<b>(1Acre) 43,560</b>	(2) 0.75 oz packets	50-100 gal	(1) 3.00 oz. packet	50-100 gal	(2) 3.00 oz. packets	100-200 gal

**NOTE:** Packets are sold in water soluble packets and re-sealable packets. Use the entire contents when using water soluble packets. Water soluble packets should NOT be used in partial amounts since they are not re-sealable. ONLY re-sealable packets of this product should be used in partial amounts.

**Low Rate:** 0.07 oz. (approx. 2 tsp) / 2-4 gal of water / 2,000 sq ft. or

**Medium Rate:** 0.07 oz. (approx. 2 tsp) / 1-2 gal of water / 1,000 sq ft.

**High Rate:** 0.14 oz. (approx. 4 tsp) / 2-4 gal of water / 1,000 sq ft

## Mixing

### **Re-sealable Mylar Packets:**

This product is sealed in mylar packets to ensure air tight and water tight seal to protect powdered product. Carefully open mylar packet and dispense the powder into the mixing tank. You can use the entire contents or you can measure a partial amount to meet mixing and rates specified for application. Airborn powder: You can reduce the amount of air born powder by avoiding windy conditions, using the entire contents of packet, and by emptying contents into a partially filled tank. Agitate tank mix.

### **Water Soluble Packets:**

Water soluble packets CANNOT be used in partial amounts. Use the entire contents of each water soluble packet. Refer to the rate chart for rate of application and for the amount of water to use for each packet. Agitate tank until entire packet has dissolved. Operate the agitator while mixing. Water soluble packets should completely dissolve within a few minutes from the time they are added to the water. Cooler water temperatures will increase the time needed for the packet to dissolve completely.

This product is packaged in a water soluble packet for your convenience. Place the packet directly into water, agitate vigorously to dissolve packet, and apply the mix solution. Do not cut water soluble packets open, they are designed to be placed directly into the tank with water. Follow label use rates for specific pests and applications.

Keep water soluble packets in a sealed container, and avoid storing packets near any moisture. Do not allow packets to become wet prior to adding into the tank. Do not handle the packets with wet hands or wet gloves. Rough handling may cause packets to break.

It's recommended to use a sticking agent to increase product coverage on plant tissue.

## CHEMIGATION

### General Information

Apply this product only through drip (trickle) or sprinkler (center pivot, lateral move, end tow, side roll, traveler, big gun, solid set, or hand move), flood (basin) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non –uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Solubilize this product with water before introduction into the system; use the diluted mixture within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH for application is a range of 5.5 to 6.5. if needed, the pH of the irrigation water can be adjusted by use of a suitable buffering agent. Agitation is necessary. Apply at the rate stated in the Directions for Use using sufficient water to achieve an even distribution within an 8 hour period. Do not apply this product at a rate that exceeds 20 grams active ingredient per acre. If applying this product in combination with other products refer to the compatibility statement in the Directions for Use section.

### **OBSERVE THE FOLLOWING PRECAUTIONS IF YOUR CHEMIGATION SYSTEM IS CONNECTED TO A PUBLIC WATER SYSTEM**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.

Chemigation systems connected to a public water systems must contain a functional, reduced-pressure zone (RPZ) , backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in the cases where there is not a water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speeds favor drift beyond the area intended for treatment.

### **STATEMENTS CONCERNING THE OPERATION OF SPRINKLER CHEMIGATION; DRIP (TRICKLE); UTILIZING A PRESSURIZED WATER AND PESTICIDE INJECTION SYSTEM**

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick – closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional , normally closed , solenoid – operated valve located on the intake side of the injection pump and connected to the system, interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

## STATEMENTS CONCERNING THE OPERATION OF FLOOD (BASIN) IRRIGATION UTILIZING GRAVITY FLOW OR PRESSURIZED WATER AND PESTICIDE INJECTION SYSTEM.

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from back flow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements.

- a. The system must contain a functional interlocking check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- b. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of the fluid back toward the injection pump.
- c. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side to the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- e. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- f. Systems must use a metering pump, such as a positive displacement injection pump, (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

### STORAGE AND DISPOSAL

**Do not contaminate water, food or feed by storage or disposal.**

**PESTICIDE STORAGE:** Do not store this product above 100°F or below 20°F for extended periods of time. Store product in the original labeled container in a cool, dry, locked place out of reach of children. Keep containers tightly closed when not in use.

**PESTICIDE DISPOSAL:** Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

## IMPORTANT: PLEASE READ BEFORE USE

### WARRANTY AND DISCLAIMER STATEMENT

To the extent consistent with applicable laws, Fortune Biotech, Ltd. warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to herein, when used in accordance with its directions; and (c) that the directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. There are no express warranties other than those set forth herein. Fortune Biotech, Ltd. neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. Fortune Biotech, Ltd. expressly excludes and disclaims all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality of performance.

This warranty does not extend to, and the user shall be solely responsible for, any loss or damage that results from the use of this product in any manner that is inconsistent with this label's directions, or cautions.

**[Optional Marketing Claims]**

- Fortune Aza 6% WP is in azadirachtin pest control.
- A way to keep insects away.
- Keep insects away with Azadirachtin
- Apply as a Spray, Soil drench, or Injection to Trees, Shrubs, Lawns and Gardens
- A water soluble product botanically derived from the neem plant, *Azadirachtin indica*.
- Fortune Aza 6% WP's solubility provides superior flow and absorption into plant tissue
- Botanical Insecticide, Repellent, Anti-feedant and Insect Growth Regulator (IGR)
- Completely water soluble and does not separate like emulsifiable concentrates (EC).

## Sub Label B

This sub-label bears direction for residential use with specific application rates associated with the package size and area to be treated.

# Fortune Aza 6% WP

This product is botanically derived from the Neem plant "Azadirachta Indica" Azadirachtin, a botanical insecticide.

### [Product Features]

- Non-oil based, water soluble powder
- Effective on a very wide spectrum of insects
- Use as a spray, drench, tree injection or chemigation
- May be applied as directed to any food crop up to and including the day of harvest

For use on turf grass, outdoor shrubs, trees and ornamentals

For ornamental greenhouse, shade house, interiors cape and nursery use

For mushroom house use

For use on outdoor food crops

For control, growth control, antifeedant and repelling Insects such as aphids, armyworms, beetles, budworms, cutworms, fungus gnats, leafhoppers, leafminers, leafrollers, lepidopterous larvae, loopers, mushroom flies, sawflies, thrips, webworms, whiteflies and other pests as listed; and plant parasitic nematodes such as dagger, golden, and root knot nematodes in vegetables, fruits, nuts, coconuts, agronomic crops and ornamental plants. For residential and commercial lawn, flowers and vegetable gardens, farms, forests, sod farms, nurseries, greenhouse-food and ornamental plants, mushrooms, nursery plants, interiorscapes, landscapes, turfgrasses, golf courses and hydroponics.

<b>ACTIVE INGREDIENT:</b> Azadirachtin .....	6.0%
<b>OTHER INGREDIENTS:</b> .....	94.0%
<b>TOTAL:</b> .....	100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>Hotline Number</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies, phone 24 hours a day, National Pesticide Telecommunication Network at 1-800-858-7378.	

See Side/Back Panel for First Aid and Precautionary Statements

**EPA Registration Number: 71038-A**

**EPA Establishment. No.: 71038-IND-1**

**Net Contents:** \_\_\_\_\_

Manufactured By:  
FORTUNE BIOTECH LIMITED  
6-6-125, Annam Gardens, Kavadiguda  
Secunderabad, 500380 India

US Contact Address:  
6860 N. Dallas Parkway, Suite 200  
Plano, TX 75024

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse.

### USER SAFETY RECOMMENDATIONS

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### PRODUCT DESCRIPTION

This product is a pale yellow/white, amorphous powder containing 6% by weight azadirachtin. It will instantly dissolve in water to give a solution that is ready for spray applications for pest control. Non-Oil based and highly effective as a powder.

Use this product for pre-harvest treatment of fruits and vegetables in case of sudden pest infestations. This product is effective on a very wide spectrum of insects and pests as listed on this label.

Use this product on a wide variety of plants as listed indoors and outdoors. If plans are made to use this product on plants not listed on this label, it is recommended that a small area such as a leaf, stem, or branch be “test sprayed”, first, and checked several days later to make sure that leaf wilting or damage does not occur.

When used as directed, this product will destroy targeted insect larvae when they, (1), eat sprayed plants, or (2), come in contact with the spray. This product eliminates insects by stopping the insect’s growth, and is effective on all insects listed, insect larval stages and pupae.

### APPLICATION INSTRUCTIONS

#### READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE

To apply this product select a suitable a hand held trigger type sprayer that will deliver a forceful, fine, leaf and fruit covering, wetting, spray mist. To get thorough spray coverage on waxy or pubescent plant surfaces the addition of small amount of a suitable sticker agent (such as NuFilm P) added to the spray mix, at the directed rates may give better foliage, insect coverage and control.

**APPLICATION METHOD AND EQUIPMENT:** Apply this product as a foliar spray to control insects and nematodes. Apply this product through sub-surface soil treatment equipment (e.g. lawn grass). To repel adult flies, apply through fogging equipment. Always follow equipment manufacturers use directions.

Apply this product by using any powered or manual pesticide application equipment. Follow the original manufacturer’s instructions when using these types of equipment.

**For optimum results, make 2 to 3 applications made at 7 to 10 day intervals, unless otherwise specified.** Foliar applications should be made to both sides of leaves. In addition, a sticker agent used as per the manufacturer’s instructions may improve product performance.

## **USE RATE DIRECTIONS** **FOR KEY PESTS BY USE SITE**

This product is intended for use on outdoor plants and food crops, mushroom houses, plants grown indoors or in greenhouses, shade cloth, interiors capes and nurseries. It can be used to control any of the following insects and nematodes.

This products label rates specify dry ounce (weight/acre (high rate) and tsp or tbsp/1000 sq. ft. (low rate). These label rates provide a high and low dose application of this product.

<b>Abbreviation &amp; Conversion Table</b>	
tsp	teaspoon
tbsp	tablespoon
A	acre
3 tsp = 1 tbsp	
1 A = 43,560 sq. ft.	

### **High Rate**

6oz/ 50 gal water / Acre (6oz. is approximately 170 tsp / 50 gal water /acre) (approximately 4 tsp / 1000 sq ft.)

### **Low Rate**

1 tsp/ 1 gal water / 1000 sq. ft. (1 tsp. is approximately 44 tsp / 44 gal water / acre)

### **COMMON USE CONDITIONS:**

**Low Rate** (Commonly used for preventative treatments before signs of infestation.)

1 tsp/ 1000 sq. ft.

**Medium Rate** (Used for most treatments. For preventative to medium infestations when pests are present.) 2-3 tsp/ 1000 sq. ft.

**High Rate** (Used for difficult to manage pests or high infestations.)

4 tsp/ 1000 sq. ft.

Use the tables below to determine the appropriate use rate for your site / pest combination.

**USE RATE DIRECTIONS**  
**FOR KEY PESTS BY USE SITE**

This product is intended for use on outdoor plants and food crops, and plants grown indoors. It can be used to control any of the following insects and nematodes.

Use the tables below to determine the appropriate use rate for your site / pest combination.

**PEST CONTROL CHART: USE RATES for indoor and outdoor plants including, FOOD CROPS, TREES and TURFGRASS.**

<b>PEST</b>	<b>RATES: Fortune Aza 6% WP oz's./Acre-tsp./1,000 sq ft.</b>	<b>COMMENTS For Spray or Drench</b>
<b>WHITEFLIES</b> , such as: Greenhouse whiteflies, Silverleaf white flies, Woolly whiteflies.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Make sure that spray covers upper, lower and all surfaces of leaves fruit and twigs.
<b>LEAF MINERS</b> , such as Azalea leafminers, Birch leafminers, Citrus leafminers, Serpentine leafminers	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Apply to new growth in spring before new larvae enter plant foliage. Repeat application at 10 to 14 day intervals if new infestations are expected.
<b>SCALE</b> , Crawlers: such as Brown Soft Scale, California red scale, Coffee Scale, Olive Scale, San Jose Scale.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft .	Make sure to thoroughly spray upper, lower and all surfaces of leaves and twigs.
<b>MEALY BUGS</b> Such as Citrus Mealybugs	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray to thoroughly cover twigs and leaves.
<b>THRIPS</b> , such as: Citrus thrips, Onion thrips, thrips palmi,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray in spring when young nymphs first appear on foliage.
<b>APHIDS</b> , such as: Cotton aphids, Green peach aphids, Pea aphids, Potato aphids	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray to wet lower side of leaves when "leaf curl" first appears.
<b>PSYLLIDS</b> , such as: Pear psylla	Same as above	Spray for new "instar" nymphs appearing on new discolored foliage.
<b>BUGS</b> , Nymphs of: such as Box-elder bugs, Chinch bugs, lygus bugs, spittle bugs, stink bugs,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray early when nymphs are young. This product will control "instar" growth until they die.
<b>FLIES</b> , Larvae of: such as Blueberry Maggot, Cherry Maggot, Crane Flies, Fruit flies, Midges, Onion Maggots, Tip worms, Walnut husk fly larvae.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	For Food and Non-food crops spray when larvae first appear.
<b>SAWFLIES</b> , Larvae of: such as: European Pine Sawflies, Yellow headed pine sawflies,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray when first larvae appear when plants start new growth.
<b>CATERPILLARS</b> , Such as: Armyworms, Artichoke plume moth, Bagworms, Bollworms, Budworms, Cabbage butterflies, Cabbage loopers, Cankerworms, Caseworms, Corn Earworms, Cutworms, Diamond back moths, Fireworms, Fruitworms, Grapeleaf skeletonizer, Gypsy moths, Hornworms, Imported Cabbage worm, leaf perforators,	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray when first larvae worms appear. Repeat applications in 7 to 10 days. For continued pest control in the spring or fall when insect infestations are expected spray ornamentals and other plants at intervals of 2 to 3 weeks.

Leafrollers, Melonworms, Navel orangeworms, Oblique banded Leafrollers, Omnivorous Leafrollers, oriental fruit moths, Pickleworms, Pine tip moths, Pinworms, Red banded leaf rollers, Sod webworms, Soybean loopers, Tent Caterpillars, Tobacco budworms, Tussocks moth larvae.		
<b>BETLES</b> , Larvae of: such as Bark beetles, Blueberry Flea beetles, Boll weevils, Colorado potato beetles, Flea beetles, Japanese beetles, Leaf beetles, Mexican, bean beetles, Phylloxera, Rose Chafers, Twig girdlers	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Spray when pests first appear. For Food Crops. Repeat application after 7 to 10 days. <u>Do not use with oil!</u> Make sure that all plant surfaces are thoroughly spray treated. Repeat in 5 to 7 days if required.
<b>WEEVILS</b> , Such as Black vine weevils, Pepper weevils, Strawberry vine weevils.	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Foliar anti-feedant sprays will stop adult feeding. Make at least 3 to 4 applications 10 days apart.
<b>BORERS</b> , Larvae such as: Peach twig borer, Peach tree borers, Cranberry borers, Emerald Ash borer (see Table below).	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	Thoroughly spray in spring after egg hatch to control young larvae.
<b>MOLE CRICKETS</b> , nymphs and young “in-stars”. <u>Turf Treatment.</u>	6 ounces in 50 gal water/A 1 tsp /1 gal. water/1,000 sq ft.	For turfgrass, spray to drench turf for young cricket nymphs in spring. Stops young from growth to adults.
<b>MUSHROOM FLIES</b> , Nematodes and Phorid Flies	Mix ½ oz. in 1 to 2 gallon of water and mist over, (or drench) 1,000 sq ft.	See “For Mushrooms” Section on this label.

**APPLICATION RATES FOR EMERALD ASH BORER**

<b>Tree Circumference in Inches</b>	<b>Emerald Ash Borer Preventative Treatment (ml)</b>	<b>*Emerald Ash Borer Symptomatic or Attacked Tree Treatment (ml)</b>
9	40	40
13	40	60
16	40	60
19	40	80
22	40	80
25	40	100
28	40	120
31	60	120
35	60	140
38	60	160
41	60	160
44	80	180
47	80	180
50	80	200
53	80	220
57	100	220
60	100	240
63	100	260
66	100	260
69	120	280
72	120	280
75	120	300
79	120	320
82	140	320
85	140	340
88	140	360
91	140	360
94	160	380
97	160	380
100	160	400
104	160	420
107	180	420
110	180	440
113	180	460
116	180	460
119	200	480
122	200	480
126	200	500
157	260	620
188	300	760

\*Emerald Ash Borer – Signs and Symptoms of Symptomatic or Attacked trees – A sign is physical damage to a tree, such as a gallery, A “D” shaped hole, or a feeding notch in the leaf, resulting from attack. A symptom is a tree’s response to attack and includes premature yellowing of foliage, dead branches, thinning crowns, or bark cracks. The effectiveness of this product in controlling Emerald Ash borer may be reduced when an ash tree has greater than 30% dead branches or thinning crown.

## USE SITES

This product can be used on Green-house: food crops, such as: Brassica (cole) crops, cucurbits, eggplants, herbs and spices, legumes, peppers, tomatoes.

**MUSHROOMS**, Varieties such as: Agaricus, enoki, maitake, oyster, shitake, and other specialty mushrooms

**FOOD CROPS**, including:

**Root and tuber vegetables**, such as: Artichoke, beets carrots, ginger, horseradish, potatoes, radishes, rutabagas, sweet potatoes, turmeric, turnips, yams.

**Leafy vegetables** (including Brassica leafy vegetables), such as: Amaranth, broccoli, Brussels sprouts, cabbage, cauliflower, celery, chervil, Chinese cabbage, collards, cress, endives, fennel, kale, kohlrabi, lettuce, mizuna ('mustard greens, parsley, purslane, rape greens, rhubarb, spinach, Swiss chard.

**Legume vegetables**, such as: beans (field, kidney etc.), chick-peas, cowpeas, guar, jackbeans, lablab beans, lentils, peas, pigeon peas, soybeans, sword beans.

**Fruiting vegetables** such as: Eggplants, ground-cherries, pepinos, peppers, pimentos, tomatillos, and tomatoes.

**Cucurbit vegetables**, such as: bitter melons, Chayotes, Chinese wax gourds, citron melons, cucumbers, gherkins, gourds, **muskmelons** (such as cantaloupes, casabas cranshaw etc.) pumpkins, squash, watermelons.

**Citrus fruits** such as: Calamondins, citrus citrons, citrus hybrids, Grapefruits, Kumquats, Lemons, Limes, Mandarins, Oranges, pumellos, satsuma mandarins.

**Pome fruits**, such as: Apples, crabapples, loquats, mayhaws, oriental pears, pears, and quinces.

**Stone fruits** such as: Apricots, cherries, nectarines, peaches, plums, and prunes.

**Berries** such as: Blackberries, caneberries, blueberries, currants, cranberries, elderberries, gooseberries, huckleberries, loganberries, raspberries, strawberries, youngberries.

**Cereal grains** such as: Barley, buckwheat, corn, millet, oats, popcorn, rice, rye, sorghum, teosinites, triticale hybrids, wheat, and wild rice.

**Herbs and spices**, including but not limited to: allspice, ahgelica, anise, annatto, balm, basil, black and white peppers, borage, burnet, camomile, caper buds, cardamom, caraway, cassia, catnip, celery seeds, chervil, chives, cinnamon, caraway, cloves, coriander (cilantro), costmary, cumin, curry leaf, dills, fennels, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, mace, marigolds, marjoram, mustard seeds, nasturtium, nutmeg, parsley, pennyroyal, pepper (black & white), poppy seeds, rosemary, rue, saffron, sage, savory, sweet bay (bay leaf), tansy, tarragon, thyme, vanilla, wintergreen, woodruff, wormwood.

**Bulb vegetables** such as: Garlic, leek, onions, shallots.

**Nuts**, such as: Almonds, beechnuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, Coconuts, filberts, hickorynuts, Imacadamia, pecans, pistachios, walnuts.

**Oilseed crops** such as: Canola, castor, crambe, guar, jojoba, peanuts, rape, safflower, sesame, soybean, sunflower.

**Tropical fruits:** such as Atemoyas, bananas, breadfruits, cherimoyas, durians, guavas, malangas, mangos, papayas, passionfruits, starfruits.

**Other food & non-food crops:** such as Asparagus, avocados, birdseed, cacao, coffee, edible flowers, feijoa, figs, ginseng, grapes, guayule, hops, kiwis, okras, olives, palms, papayas, pawpaws, persimmons, pineapples, rambutans, sugarcane, tamarillos, tea, tobacco, water chestnuts, watercress.

**Ornamental Plants:** such as: African violets, ageratum, aster, aucuba, begonia, cacti, calendula, calla, carnation, ceanothus, chrysanthemum, cineraria, coleus, cyclamen, daffodil, dahlia, delphinium, ficus, foliage plants, fuschia, gardenia, geranium, gloxinia, hyacinth, hydrangea, iris, ivy, lily, maidenhair fern, marigold, narcissus, orchid, pansy, pelargonium, peony, phlox, pittisporum, poinsettia, pyracantha, rubber plant, snapdragon, stock, tulip, wandering jew, yew, yucca, zinnia.

**Ornamental Trees and Shrubs:** such as: Andromeda, Arbovitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, camellia, cedar, chamaecyprus, dogwood, douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honeylocust, horsechestnut, ilex, juniper, larch laurel, lilac, linden, London plane, magnolia, manvillia, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, pine, phinota, pines, plane tree, poplar, privet, quince, rhododendron, roses, spruce, sycamore, white cedar and white pine.

### FOR TURF GRASSES

Bent grass	Fescue
Bermuda grass	Ryegrass; annual
Bluegrass annual & perennial	Ryegrass; perennial
Buffalo grass	Wheat grass
Centipede grass	Zoysia grass

For control of Sod Webworms, Cutworms, Aphids, Leafhoppers, ants, and chiggers: use a suitable pressure sprayer and mix 1 tbsp. in 2 to 3 gal. of water and apply to 2,500 sq ft of turf. Apply when insect larvae first appear and if necessary repeat application in 10 to 14 days. The use of an approved “spreader sticker” may help the spray to penetrate turf down to the larvae/worm feeding area.

Packet Size	Mylar Packet	Water Soluble Packet
0.07 oz. packets (approx. 2tsp)	not available	yes
0.75 oz. packets (approx. 22tsp)	yes	yes
3.00 oz. packets (approx. 85tsp)	yes	yes

## Dose/Rate

### Spray Applications

This product is measured in dry ounces (weight) and approximate teaspoons for each packet size. Packets come in various sizes. Depending on the type of pest and timing of treatment you may use low, medium, or high rates of application.

**NOTE: Low Rates** are commonly used for preventative treatments before signs of insects. **Medium Rates** are used for most treatments for preventative to medium infestations when pests are present. **High Rates** are used for difficult to manage pests or for heavy infestations.

### *Product packet directions for mixing, and dosing in spray applications.*

*Packet sizes are determined by using the most economical size and no partial packets.*

Sq Ft.	Low Rate		Medium Rate		High Rate	
	Product	Water	Product	Water	Product	Water
<b>1,000</b>			(1) 2tsp packet	1-2 gal	(2) 2tsp packets	1-4 gal
<b>5,000</b>	(3) 2tsp packets	5-10 gal	(5) 2tsp packets	5-10 gal	(1) 0.75 oz packet	10-20 gal
<b>10,000</b>	(5) 2tsp packets	10-20 gal	(1) 0.75oz packet	10-20 gal	(2) 0.75 oz packets	20-40 gal
<b>20,000</b>	(1) 0.75oz packet	20-40 gal	(2) 0.75 oz packets	20-40 gal	(1) 3.00 oz. packet	50-100 gal
<b>(1Acre) 43,560</b>	(2) 0.75 oz packets	50-100 gal	(1) 3.00 oz. packet	50-100 gal	(2) 3.00 oz. packets	100-200 gal

**NOTE:** Packets are sold in water soluble packets and re-sealable packets. Use the entire contents when using water soluble packets. Water soluble packets should NOT be used in partial amounts since they are not re-sealable. ONLY re-sealable packets of this product should be used in partial amounts.

**Low Rate:** 0.07 oz. (approx. 2 tsp) / 2-4 gal of water / 2,000 sq ft. or

**Medium Rate:** 0.07 oz. (approx. 2 tsp) / 1-2 gal of water / 1,000 sq ft.

**High Rate:** 0.14 oz. (approx. 4 tsp) / 2-4 gal of water / 1,000 sq ft.

## Mixing

### Re-sealable Mylar Packets:

This product is sealed in mylar packets to ensure air tight and water tight seal to protect powdered product. Carefully open mylar packet and dispense the powder into the mixing tank. You can use the entire contents or you can measure a partial amount to meet mixing and rates specified for application. Airborne powder: You can reduce the amount of air born powder by avoiding windy conditions, using the entire contents of packet, and by emptying contents into a partially filled tank. Agitate tank mix.

### Water Soluble Packets:

Water soluble packets CANNOT be used in partial amounts. Use the entire contents of each water soluble packet. Refer to the rate chart for rate of application and for the amount of water to use for each packet. Agitate tank until entire packet has dissolved.

Operate the agitator while mixing. Water soluble packets should completely dissolve within a few minutes from the time they are added to the water. Cooler water temperatures will increase the time needed for the packet to dissolve completely.

## Retail Supplement

This product is packaged in a water soluble packet for your convenience. Place the packet directly into water, agitate to dissolve packet, and apply the mix solution.

### Net Contents:

0.28 oz. (approx. 8 tsp) of product

4 water soluble packets 0.07 oz. each (approx. 2 tsp each)

\* Full contents treats up to 8,000 sq ft. for spray application (2-8 gallons)

\* Full contents treats 1 medium size tree [dbh] for soil drench application (2 gallons)

**NOTE: Low Rates** are commonly used for preventative treatments before signs of insects. **Medium Rates** are used for most treatments for preventative to medium infestations when pests are present. **High Rates** are used for difficult to manage pests or for heavy infestations.

### ***Retail Directions: (0.07oz.) 2 teaspoon packet mixing and dosing for spray applications***

2 tsp packets	Low Rate		Medium Rate		High Rate		
	Sq Ft.	Product	Water	Product	Water	Product	Water
1,000				(1) 2tsp packet	1-2 gal	(2) 2tsp packets	1-4 gal
2,000	(1) 2tsp packets		2-4 gal	(2) 2tsp packets	2-4 gal	(4) 2tsp packets	2-8 gal
5,000	(3) 2tsp packets		5-10 gal	(5) 2tsp packets	5-10 gal	(10) 2tsp packets	10-20 gal
10,000	(5) 2tsp packets		10-20 gal	(10) 2tsp packets	10-20 gal	(20) 2tsp packets	20-40 gal

**NOTE:** Packets are sold in water soluble packets. Use the entire contents when using water soluble packets. Water soluble packets should NOT be used in partial amounts since they are not re-sealable.

**Low Rate:** 0.07 oz. (approx. 2 tsp) / 2-4 gal of water / 2,000 sq ft. or

**Medium Rate:** 0.07 oz. (approx. 2 tsp) / 1-2 gal of water / 1,000 sq ft.

**High Rate:** 0.14 oz. (approx. 4 tsp) / 2-4 gal of water / 1,000 sq ft.

### Soil Applications to Trees

Soil Drench	Pest	Comments
<b>Small tree (2-7" DBH):</b> 4 tsp / gal of water  <b>Medium tree (8-15" DBH):</b> 8-10 tsp / gal of water  <b>Large tree (16"+ DBH):</b> 16-22 tsp / gal of water	Ips engraver beetles Flatheaded borers (e.g., bronze birch borer) Leaf miners (e.g., birch leafminer)	Remove organic matter from the base of the tree, apply to soil around the tree base, 1 - 3 feet from trunk or inject (6" deep) into the soil working around the base of the tree.  Make the first application to trees 1-2 weeks prior to expected adult emergence. For pests with multiple generations, repeat applications once every 3 to 4 weeks
	Leaf defoliating insects (including winter moth, fall webworm)	Remove organic matter from the base of the tree, apply to soil around the tree base, 1 - 3 feet from trunk or inject (6" deep) into the soil working around the base of the tree.  Make application when caterpillars first hatch; prior to bud break.

### Bark Spray Applications to Trees

Bark Spray	Pest	Comments
<b>Small tree (2-7" DBH):</b> 2 tsp / gal of water	Conifer bark beetles (e.g., black turpentine beetle, mountain pine beetle)	Spray the bole of tree thoroughly covering the bark from the ground up 20 feet.
<b>Medium tree (8-15" DBH):</b> 4-5 tsp / gal of water	Flatheaded borers (e.g., bronze birch borer)	Make the first application to trees 1-2 weeks prior to expected adult emergence. For pests with multiple generations, repeat applications once every 10 to 14 days
<b>Large tree (16"+ DBH):</b> 8-11 tsp / gal of water		

### Applications to Lawns

Pest	Soil Drench	Comments
Annual Bluegrass Weevil	4-8 tsp / 2-3gal of water / 1,000 sq ft.	Make applications in early spring (in the northeast, 3 <sup>rd</sup> or 4 <sup>th</sup> week of April) when weevils first become active
White Grubs	4-8 tsp / 2-3gal of water / 1,000 sq ft.	Make applications in early spring and again in summer when larvae are feeding

### Water Soluble Packet Supplement

This product is packaged in a water soluble packet for your convenience. Place the packet directly into water, agitate vigorously to dissolve packet, and apply the mix solution. Do not cut water soluble packets open, they are designed to be placed directly into the tank with water. Follow label use rates for specific pests and applications.

Keep water soluble packets in a sealed container, and avoid storing packets near any moisture. Do not allow packets to become wet prior to adding into the tank. Do not handle the packets with wet hands or wet gloves. Rough handling may cause packets to break.

#### STORAGE AND DISPOSAL

**Do not contaminate water, food or feed by storage or disposal.**

**PESTICIDE STORAGE:** Store product in the original labeled container in a cool, dry, locked place out of reach of children. Keep containers tightly closed when not in use.

**PESTICIDE PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the container be used, carefully following label directions and precautions.

**CONTAINER HANDLING; IF EMPTY:** Non-refillable container. Do not reuse this container. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or for disposal instructions. Never place unused product down any indoor or outdoor drain.

### IMPORTANT: PLEASE READ BEFORE USE

#### WARRANTY AND DISCLAIMER STATEMENT

To the extent consistent with applicable laws, Fortune Biotech, Ltd. warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to herein, when used in accordance with its directions; and (c) that the directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. There are no express warranties other than those set forth herein. Fortune Biotech, Ltd. neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. Fortune Biotech, Ltd. expressly excludes and disclaims all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality of performance.

This warranty does not extend to, and the user shall be solely responsible for, any loss or damage that result from the use of this product in any manner that is inconsistent with this label's directions, or cautions.

**[Optional Marketing Claims]**

- Fortune Aza 6% WP is in azadirachtin pest control.
- A way to keep insects away.
- Keep insects away with Azadirachtin
- Apply as a Spray, Soil drench, or Injection to Trees, Shrubs, Lawns and Gardens
- A water soluble product botanically derived from the neem plant, *Azadirachtin indica*.
- Fortune Aza 6% WP's solubility provides superior flow and absorption into plant tissue
- Botanical Insecticide, Repellent, Anti-feedant and Insect Growth Regulator (IGR)
- Completely water soluble and does not separate like emulsifiable concentrates (EC).